

β -Actin Mouse mAb

Catalog No.: AC004 **282 Publications**

Basic Information

Observed MW

42kDa

Calculated MW

42kDa

Category

Loading control antibody

Applications

ELISA, WB, IHC-P, IF/ICC

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

AMC0001

Recommended Dilutions

WB	1:2000 - 1:20000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

Contact

 | www.abclonal.com

Background

This gene encodes one of six different actin proteins. Actins are highly conserved proteins that are involved in cell motility, structure, integrity, and intercellular signaling. The encoded protein is a major constituent of the contractile apparatus and one of the two nonmuscle cytoskeletal actins that are ubiquitously expressed. Mutations in this gene cause Baraitser-Winter syndrome 1, which is characterized by intellectual disability with a distinctive facial appearance in human patients. Numerous pseudogenes of this gene have been identified throughout the human genome.

Immunogen Information

Gene ID	Swiss Prot
60	P60709

ImmunogenFusion protein of human β -Actin**Synonyms**

BRWS1; PS1TP5BP1

Product Information


Source	Isotype	Purification
Mouse	IgG1,Kappa	Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of extracts of HeLa cells, using β -Actin Mouse mAb (AC004) at 1:2000- 1:640000 dilution.


Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.

Lysates/proteins: 25 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.



Western blot analysis of extracts of various lysates, using β -Actin Mouse mAb (AC004) at 1:20000 dilution.


Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.

Lysates/proteins: 25 μ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Immunofluorescence analysis of U2OS cells using β -Actin Mouse mAb (AC004) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.